

# Interactions of the Effects of Provenances and Habitats on the Growth of Scots Pine in Two Provenance Tests in Bosnia and Herzegovina

Mirzeta Memišević Hodžić<sup>1,\*</sup>, Dalibor Ballian<sup>1,2</sup>

(1) University of Sarajevo, Faculty of Forestry, Department of Silviculture and Urban Greenery, Zagrebačka 20, BA-71000 Sarajevo, Bosnia and Herzegovina; (2) Slovenian Forestry Institute, Večna pot 2, SI-1000 Ljubljana, Slovenia

\* Correspondence: e-mail: [m.memisevic-hodzic@sfsa.unsa.ba](mailto:m.memisevic-hodzic@sfsa.unsa.ba)

**Citation:** Memišević Hodžić M, Ballian D, 2021. Interactions of the Effects of Provenances and Habitats on the Growth of Scots Pine in Two Provenance Tests in Bosnia and Herzegovina. *South-east Eur for* 12(1): early view. <https://doi.org/10.15177/seefor.21-03>.

**Received:** 11 Jan 2021; **Revised:** 9 Mar 2021; **Accepted:** 12 Mar 2021; **Published online:** 11 Apr 2021

## ABSTRACT

This research aims to determine the interaction of the effects of provenance and habitat conditions on provenance tests on the growth of Scots pine on two experimental plots in Bosnia and Herzegovina. Provenance tests are located on plots with different ecological conditions and altitudes: Romanija Glasinac, 1000 m, and Gostović Zavidovići, 480 m. Both tests include 11 provenances and two clonal seed plantations with 10 families in each, and five repetitions. Tree heights and diameters at breast height were measured at the age of 21 years. Interactions were determined using multivariate analysis for measured traits. The highest average heights on the provenance test Glasinac had provenances Bugojno, Romanija Glasinac, and Šipovo (8.8 m), and on the Gostović provenance Rogatica (11.0 m). The highest average diameter at breast height on the Glasinac test had Šipovo provenance (13.9 cm) and on the Gostović test Bosanski Petrovac provenance (12.3 cm). Variance analysis showed statistically significant differences among provenances in terms of diameter at breast height and height values. Multivariate analysis showed the presence of interactions of effects of provenances and habitat conditions on provenance tests. For provenances that did not show interaction, it is recommended to use provenances that performed better in the given ecological conditions, and for those that showed interaction, it is necessary to choose those provenances which are expected to show better results in given conditions later in life. The obtained results are very important for the conservation activities of this species.

**Keywords:** *Pinus sylvestris*; morphological traits; genetic structure; environmental conditions

## INTRODUCTION

Scots pine (*Pinus sylvestris* L.) is one of the most important tree species in European forests. It is characterized by high genetic variability and complex population structure (Stefanović et al. 1980, Omanović 2008). In Bosnia and Herzegovina, differences within Scots pine populations can be expected at the morphological level (Stefanović 1980). Differences can be expected in the production potential of provenances as well (Mikić 1991). Due to the importance of this species for the forestry of Bosnia and Herzegovina, clonal seed plantations were established (Ballian and Božić 2004, Ballian et al. 2005), two of which were used in this research.

Knowing whether a species has resilience and adaptability to changing environmental conditions, as well as stress conditions, is important for the functioning of modern forestry and conservation of forest ecosystems. Higher genetic variability makes the species more resilient and better adapted for survival in changing environmental conditions and in stress conditions (Villani and Eriksson 2006).

There have not been many studies on the interactions between the effects of provenance genetic structure and habitat conditions on Scots pine. Some authors studied morphological traits of Scots pine in international provenance tests in Poland (Barzdajn et al. 2016), Spain (Alía et al. 2001), Turkey (Gülcü and Bilir 2017), and in

Bosnia and Herzegovina (Ballian et al. 2009, Cvjetković et al. 2014, Ballian and Šito 2017, Ballian et al. 2019). Taeger et al. (2013) studied the impact of climate and drought events on the growth of Scots pine provenances in an international provenance test which also contained a provenance from Bosnia and Herzegovina, and the results indicate a better adaptation to drought of the provenance from Bosnia Herzegovina and of local German provenances compared to other provenances.

Unfortunately, so far there has been little research dealing with provenance interactions. So far, interactions among provenances originating from different countries, from a greater geographical distance (Memišević Hodžić et al. 2020), but not provenances from a narrower geographical area such as Bosnia and Herzegovina, have been analyzed. Thus, Memišević Hodžić et al. (2020) studied the interaction between the effects of provenance genetic structure and habitat conditions on international provenance tests of Scots pine in two locations in Bosnia and Herzegovina. Obtained results indicated interactions on some provenances.

For studying the variability and adaptability of Scots pine in Bosnia and Herzegovina, provenance tests were established in 1988 on two locations, Romanija Glasinac and Zavidovići Gostović. The study aims to determine whether there is an interaction between the effects of provenance genetic structure and habitat conditions on the growth of Scots pine in both test plots, for selection and breeding of provenances adapted to habitat conditions.

## MATERIALS AND METHODS

In this paper, the material from two provenance tests of Scots pine in Bosnia and Herzegovina was investigated. Information about the locations of test plots are shown in Table 1.

Provenance tests were established from seed material from 11 seed stands of Scots pine and two clonal seed plantations (Sarajevo-Rakovica and Dobož-Stanovi). A random block system was applied, and the distribution of provenances was the same at both provenance tests. The seeds were sown in 1986 in the nursery "Slatina" of the Faculty of Forestry in Sarajevo. At the age of two years (1 + 1), seedlings were planted in two experimental plots, with planting spacing of 2x2m. Each provenance within the plot was composed of ten families with four plants, and each provenance was planted in five replications. The list of provenances and information about their locations is shown in table 2 and their geographical distribution in figure 1.

In the spring of 2007, plant heights and diameters at breast height were measured. Data were processed using the statistical program SPSS 26.0. Descriptive and multivariate analyses (Two-way ANOVA) were performed. Two-way ANOVA shows whether there is an interaction of two independent factors (provenance and habitat) on the dependent variable (height and diameter). In other words, it shows whether the influence of one of the independent variables on the dependent variable is the same for all values of the second independent variable (and vice versa).

## RESULTS

Average values of height per provenances and provenance tests are shown in Figure 2. All provenances had higher values of average heights on the Gostović than on the Glasinac plot. The highest average heights on the Glasinac plot had provenances Bugojno, Romanija Glasinac, and Šipovo (8.8 m), and on the Gostović plot provenance Rogatica (11.0 m). The lowest average height on the Glasinac plot had Olovo and Zavidovići provenances (7.3 m), and on the Zavidovići plot seed plantation Dobož had the lowest average height (9.0 m).

**Table 1.** Information about Scots pine provenance tests' locations.

Information / Provenance test	Romanija Glasinac	Zavidovići Gostović
North Latitude	43°52'43.10"	44°20'55.27"
East Longitude	18°50'31.34"	18°08'04.03"
Forest Management Area	„Sokolačko“	„Krivajsko“
Management Unit	„Romanija Glasinac“	„Gostović“
Department	15	42
Altitude	1000 m	480 m
Exposition	E	SW
Inclination	5°	20°
Geological substrate	Limestone	Serpentine-peridotite
Soil type	Calcomelanosol	Ranker, eutric cambisol
Vegetation	<i>Piceo Pinetum illyricum pyroletosum</i>	<i>Pinetum silvestris nigrea serpentinicum</i>
Average annual temperature	7.08°C	9.35°C
Vegetation period duration	157 days	184 days

**Table 2.** List and location information of provenances in the provenance test.

Provenance	North Latitude	East Longitude	Altitude (m)	Soil type	Geological substrate
Bosanski Petrovac	44°29'28"	16°29'21"	920	rendzina	dolomite, limestone
Bugojno	44°03'10"	17°19'55"	1230	dolomite rendzina	dolomite
Foča	43°20'51"	18°56'25"	1370	distric cambisol	quartzporphyrite
Han Kram	44°01'53"	18°56'32"	1100	swampy, charred, brown podzolic	tuff sandstones
Kladanj	44°16'46"	18°37'47"	900	district cambisol and brown podzolic soil	sandstones, clay minerals, cherts
Olovo	44°14'50"	18°20'54"	850	eutric cambisol	peridotite
Rogatica	43°48'06"	19°08'32"	1100	distric cambisol, luvisol	cherts
Romanija – Glasinac	43°54'20"	18°42'50"	1235	calcocambisol	limestone
Romanija – Palež	44°02'14"	18°45'50"	780	pseudogley	diabase gabbro
Seed plantation Doboj	44°45'11"	17°59'50"	180	fluvisol	sandstone
Seed plantation Sarajevo	43°52'14"	18°12'52"	570	clay, ortstein	marl
Šipovo	44°14'31"	17°13'26"	1200	dolomite rendzina, calcocambisol	dolomite, limestone
Zavidovići	44°19'56"	18°12'44"	480	ranker, eutric cambisol	serpentine peridotite

**Figure 1.** Provenances included in provenance tests Glasinac and Gostović (SP – seed plantation; PT – provenance test).

Average values of breast height diameter per provenances are shown in Figure 3. Unlike for the trait of height, all provenances had higher average values of diameter at breast height on the Glasinac than on the Gostović plot. The highest average breast diameter on the Glasinac plot had Šipovo provenance (13.9 cm) and on the Gostović plot Bosanski Petrovac provenance (12.3 cm). The lowest average values of breast height diameter on the Glasinac plot had provenance Olovo (11.9 cm) and on the Gostović plot seed plantation Doboj (9.8 cm).

In the multivariate tests table for height and diameter at breast height together (Table 3) it is visible that Wilks' Lambda showed Sig. value of 0.000, which indicates  $p <$

0.0005 for both factors and interaction between factors. It means that values of heights and diameters at breast height are significantly dependent on provenances, habitat conditions on provenance tests, and interaction of provenances and habitat conditions on provenance tests.

Considering that an interaction effect can usually be seen as a set of non-parallel lines, it is visible in Figure 4 that there is a statistically significant interaction of effects of provenance and environmental conditions on experimental plots for the trait of height for some of the researched provenances. The provenance of seed plantation Doboj was most clearly distinguished by the interaction of provenance x habitat, with estimated marginal mean of height among




Figure 2. Average values of height of plants per provenances and experimental plots.




Figure 3. Average values of breast height diameter of plants per provenances and tests.

other provenances on Glatinac, but lower than other provenances on the Gostović plot. Provenance Romanija Glasinac had one of the highest estimated marginal means on the Glatinac plot, but one of the lowest marginal means on the Gostović plot. Provenance Rogatica had estimated marginal mean of height about average for all provenances on the Glatinac plot, but the highest estimated marginal mean on the Gostović plot. Provenance Šipovo had the highest estimated marginal mean for height on the Glatinac plot, and one of the highest marginal means on the Gostović


plot. Provenances Zavidovići and Olovo had the lowest estimated marginal means for height on the Glatinac plot, but on the Gostović plot, their values were among the values of other provenances.

Tests of between-subjects effects table (Table 4) for the trait of height showed the presence of statistically significant interaction between provenances and habitat conditions (Fizr.>Ftab., Sig.<0.005).

In the graphical representation of estimated marginal means for the trait of diameter at breast height (Figure 5),

**Table 3.** Results of multivariate tests for the height and diameter at breast height.

	Effect	Value	F	Hypothesis df	Error df	Sig.
Intercept	Pillai's Trace	0.961	45238.595 <sup>b</sup>	2.000	3651.000	0.000
	Wilks' Lambda	0.039	45238.595 <sup>b</sup>	2.000	3651.000	0.000
	Hotelling's Trace	24.781	45238.595 <sup>b</sup>	2.000	3651.000	0.000
	Roy's Largest Root	24.781	45238.595 <sup>b</sup>	2.000	3651.000	0.000
Provenance test	Pillai's Trace	0.541	2150.029 <sup>b</sup>	2.000	3651.000	0.000
	Wilks' Lambda	0.459	2150.029 <sup>b</sup>	2.000	3651.000	0.000
	Hotelling's Trace	1.178	2150.029 <sup>b</sup>	2.000	3651.000	0.000
	Roy's Largest Root	1.178	2150.029 <sup>b</sup>	2.000	3651.000	0.000
Provenance	Pillai's Trace	0.030	4.656	24.000	7304.000	0.000
	Wilks' Lambda	0.970	4.669 <sup>b</sup>	24.000	7302.000	0.000
	Hotelling's Trace	0.031	4.682	24.000	7300.000	0.000
	Roy's Largest Root	0.025	7.655 <sup>c</sup>	12.000	3652.000	0.000
p. test * Provenance	Pillai's Trace	0.030	4.690	24.000	7304.000	0.000
	Wilks' Lambda	0.970	4.713 <sup>b</sup>	24.000	7302.000	0.000
	Hotelling's Trace	0.031	4.736	24.000	7300.000	0.000
	Roy's Largest Root	0.028	8.602 <sup>c</sup>	12.000	3652.000	0.000

**Figure 4.** Graph of the interaction of effects of provenance and habitat conditions for height

it is visible that there is a statistically significant interaction between provenance and environmental conditions of experimental plots for some of the researched provenances. Similar to the trait of height, the provenance of seed plantation Dobož was most clearly distinguished by the interaction of provenance x habitat for the breast height diameter, with estimated marginal mean among other provenances on Glasinac, but lower than other provenances on the Gostović plot. Provenance Romanija Glasinac had one of the highest estimated marginal means on the Glasinac plot, but one of the lowest marginal means on the Gostović plot. Provenance Rogatica had estimated marginal mean of diameter at breast height about average for all

provenances on the Glasinac plot, but the highest estimated marginal mean on the Gostović plot. Provenance Šipovo had the highest estimated marginal mean for diameter at breast height on the Glasinac plot, and one of the highest marginal means on the Gostović plot. Provenance Olovo had the lowest estimated marginal mean for diameter at breast height on the Glasinac plot, but on the Gostović plot, its value was among the values of other provenances.

Tests of between-subjects effects table (Table 5) for the trait of breast height diameter showed that there is a statistically significant interaction of effects of provenances and habitat conditions on provenance tests conditions (Fizr.>Ftab., Sig.<0.005).




Figure 5. Graph of the interaction of effects of provenance and habitat conditions for diameter at breast height.

Table 4. Tests of between-subjects effects for trait of height.

Source	Type III Sum of Squares	df	Mean Square	F	Sig.
Corrected Model	4796.276 <sup>a</sup>	25	191.851	54.368	0.000
Intercept	318609.332	1	318609.332	90290.226	0.000
Provenance test	4015.689	1	4015.689	1138.000	0.000
Provenance	311.826	12	25.986	7.364	0.000
P. test * Provenance	358.017	12	29.835	8.455	0.000
Error	12886.902	3652	3.529		
Total	342382.930	3678			
Corrected Total	17683.178	3677			

R Squared = .271 (Adjusted R Squared = 0.266)

Table 5. Tests of between-subjects effects for diameter at breast height.

Source	Type III Sum of Squares	df	Mean Square	F	Sig.
Corrected Model	2970.403 <sup>b</sup>	25	118.816	11.473	0.000
Intercept	552301.877	1	552301.877	53329.602	0.000
Provenance test	1824.163	1	1824.163	176.139	0.000
Provenance	768.520	12	64.043	6.184	0.000
P. test * Provenance	496.316	12	41.360	3.994	0.000
Error	37821.517	3652	10.356		
Total	599746.103	3678			
Corrected Total	40791.920	3677			

R Squared = .073 (Adjusted R Squared = 0.066)

## DISCUSSION

Investigating the interactions of the effects of provenances and environmental conditions can be important in the breeding strategy of Scots pine.

In this research, statistically significant interactions of effects of provenance and environmental conditions on experimental plots for both traits and some of the researched provenances were found. Provenance Romanija Glasinac had one of the highest estimated marginal

means for height on the Glasinac plot, but one of the lowest marginal means on the Gostović plot. It could be explained by the vicinity of the provenance origin to the provenance test and similar ecological conditions. Provenances Zavidovići and Olovo had the lowest estimated marginal means for height on the Glasinac plot, but on the Gostović plot, their values were among the values of other provenances, which can also be explained by the vicinity of the provenances and similarity of their ecological conditions with the provenance test Gostović. Similar to the trait of height, provenance Romanija Glasinac had one of the highest estimated marginal means for diameter at breast height on the Glasinac plot, but one of the lowest marginal means on the Gostović plot. Provenance Rogatica had estimated marginal mean of diameter at breast height about average for all provenances on the Glasinac plot, but the highest estimated marginal mean on the Gostović plot. Provenance Olovo had the lowest estimated marginal mean for height on the Glasinac plot, but on the Gostović plot, its value was among the values of other provenances.

These results correspond to the results of other researchers. Eiche and Andersson (1974) investigated survival and growth in Scots pine in provenance tests in Northern Sweden. Their results showed the geneecological variation of climatic hardiness, capacity for survival, and growth rate of different populations. They concluded that interaction for genotype and environment emerged quite obviously. Matheson and Raymond (1984, 1986) stated that the provenance by site interactions in forest trees are often detectable, but of small importance. Savva and Vaganov (2006) assessed genetic and environmental effects in Scots pine provenances planted in Central Siberia. They found that the main climatic factors controlling tree-rings formation differed slightly among different provenances. The genetically fixed ability of the provenances was not great (less than 15%), which proves the high adaptability of Scots pine to abrupt climatic change. The tree-ring formation of Scots pine provenances is mainly determined by environmental factors. Zhelev and Lust (1999) investigated Scots pine provenances in Belgium and found that the provenance by site interactions was of importance only for height growth. Alía et al. (2001) investigated 16 Spanish and 6 German provenances of Scots pine in a provenance test in Spain. Based on data for the height, diameter, the number of twigs at the fourth year whorl, and survival, provenance by site interaction was very significant ( $P < 0.01$ ) for most traits.

Ballian et al. (2009) measured the diameter and height of the plants and calculated basal area and volume of plants in the provenance test of Scots pine Glasinac Sokolac, which is one of the provenance tests in this research. The provenance geographically closest to the provenance test showed the best results of productivity. Cvjetković et al. (2014) measured and calculated the same characteristics at the same plot in 2011. The results confirmed the existence of significant differences within and among provenances, and the provenance geographically closest to the provenance test showed the best results. In this research, provenance Romanija Glasinac, closest to the provenance test Glasinac Sokolac, showed the highest average height, but not the highest average of breast height diameter.

Gülcü and Bilir (2017) studied provenance x site interactions in thirteen-year-old Scots pine provenance tests at two sites in the southern part of Turkey, containing 30 provenances, based on height, diameter, and survival of plants. The results showed statistically significant provenance x site interaction ( $p < 0.05$ ). Memišević Hodžić et al. (2020) studied provenance x site interaction in two international provenance tests in Bosnia and Herzegovina, based on the data of heights and root collar diameter, and determined interaction between some of the researched provenances.

## CONCLUSIONS

The results of this research showed interaction of effects of provenance genetic structure and habitat conditions for both researched traits (height and diameter at breast height). Considering the height, provenances Bugojno, Romanija Glasinac, and Šipovo had the highest, and provenances Olovo and Zavidovići the lowest values on the Glasinac provenance test. On the Gostović test, provenance Rogatica had the highest, and Seed plantation Doboj the lowest value of height. The highest values of breast diameter on the Glasinac provenance test had Šipovo provenance and the lowest Olovo provenance. On the Gostović test, provenance Bosanski Petrovac had the highest values of breast height diameter, and the lowest had Seed plantation Doboj. Except for provenance Romanija Glasinac on the Glasinac test for the trait of height, provenances originated from locations nearest to the provenance test locations did not show higher values of researched traits. Wilks' Lambda multivariate tests showed that values of heights and diameters at breast height are significantly dependent on provenance, habitat conditions on provenance tests locations, and interaction of provenance x habitat conditions on provenance tests locations. Graphical representation of estimated marginal means for both researched traits showed the presence of statistically significant interaction of effects of provenance and environmental conditions on provenance tests for some of the researched provenances. Tests of between-subjects effects for both traits showed a statistically significant interaction of effects of provenances and habitat conditions on provenance tests. The results of this study can be used in the process of Scots pine breeding and for its conservation by *in situ* and *ex-situ* methods.

## Author Contributions

MMH, DB, conceived and designed the research, DB carried out the field measurements, DB performed measuring, MMH processed the data and performed the statistical analysis, DB supervised the research and helped to draft the manuscript, MMH wrote the manuscript.

## Funding

This research received no external funding.

## Conflicts of Interest

The authors declare no conflict of interest.

## REFERENCES

- Alía R, Moro-Serrano J, Notivol E, 2001. Genetic variability of Scots pine (*Pinus sylvestris*) provenances in Spain: growth traits and survival. *Silva Fenn* 35(1): 27-38. <https://doi.org/10.14214/sf.601>.
- Ballian D, Bogunić F, Konner M, 2005. Usporedba molekularno genetičkih svojstava sjemenskih plantaža običnog bora (*Pinus sylvestris* L.) u Bosni i Hercegovini. *Rad Šumar Ins. Jastrebarsko* 41(2): 7-16.
- Ballian D, Božić G, 2004. Kontrola morfološke identifikacije klonova iz sjemenske plantaže bijelog bora "Koziji grm" pomoću izoenzimskih markera. *Radovi Šum Fak Sarajevo* 1: 47-56.
- Ballian D, Lizdo E, Bogunić F, 2019. Analiza diferenciranosti rasta i fenologije provenijencija običnog bora (*Pinus sylvestris* L.) u pokusu provenijencija kod Kupresa (Bosna i Hercegovina). *Sumar List* 143(1-2): 25-34. <https://doi.org/10.31298/sl.143.1-2.3>.
- Ballian D, Mujanović E, Čabaravdić A, 2009. Varijabilnosti običnog bora (*Pinus sylvestris* L.) u pokusu provenijencija Glasinac – Sokolac, Bosna i Hercegovina. *Sumar List* 133(11-12): 577-588.
- Ballian D, Šito S, 2017. Analysis of differences of growth and phenology of provenances of Scots pine (*Pinus sylvestris*) in provenance experiment at Žepče. *Biosyst Divers* 25(3): 228-235. <https://doi.org/10.15421/011735>.
- Barzdajn W, Kowalkowski W, Chmura DJ, 2016. Variation in growth and survival among European provenances of *Pinus sylvestris* in a 30-year-old experiment. *Dendrobiology* 75: 67-77. <https://doi.org/10.12657/denbio.075.007>.
- Cvijetković B, Mataruga M, Dukić V, Daničić V, Lučić A, 2014. The variability of Scots pine (*Pinus sylvestris* L.) in the provenance test in Bosnia and Herzegovina. In: Proceedings of the Biennial International Symposium Forest and Sustainable Development, Brasov, Romania, 24-25 October 2014. Transilvania University Press, Brasov, Romania, pp 79-88.
- Giertych M, Matyas C, 1991. Genetics of Scots pine. Developments of Plant Genetics and Breeding. Vol 3. Elsevier, Amsterdam, Netherlands, 280 p.
- Eiche V, Andersson E, 1974. Survival and Growth in Scots Pine (*Pinus sylvestris* L.) Provenance Experiments in Northern Sweden. *Theoret Appl Genetics* 44: 49-57. <https://doi.org/10.1007/BF00277952>.
- Gülcü S, Bilir N, 2017. Growth and Survival Variation among Scots Pine (*Pinus sylvestris* L.) Provenances. *Int J Genomics*. <https://doi.org/10.1155/2017/1904623>.
- Matheson AC, Raymond CA, 1984. Provenance x environment interaction: its detection, practical importance and use with particular reference to tropical forestry. In: Barnes RD, Gibson GL (eds) Commonwealth Forestry Institute, Oxford, and Forest Research Center, Harare, pp 81-117.
- Matheson AC, Raymond CA, 1986. A review of provenance x environment interaction: its practical use with particular reference to the tropics. *Commonw Forest Rev* 65(4): 283-302.
- Memišević Hodžić M, Bejčić S, Ballian D, 2020. Interaction Between the Effects of Provenance Genetic Structure and Habitat Conditions on Growth of Scots Pine in International Provenance Tests in Bosnia and Herzegovina. *South-east Eur for* 11(1): 11-16. <https://doi.org/10.15177/seefor.20.03>.
- Mikić T, 1991. Primjena metoda oplemenjivanja u podizanju intezivnih kultura šumskog drveća u cilju povećanja proizvodnje drvne mase sa kratkim produkcionim periodom. Izvještaj za period 1989-1990 u okviru D.C.VII. Sarajevo.
- Omanović M, 2008. Biohemijska karakterizacija prirodnih populacija običnog bora (*Pinus sylvestris* L.) u dijelu rasprostranjenja u Bosni i Hercegovini. MSc Thesis, University of Sarajevo, Faculty of Forestry Sarajevo, Bosnia and Herzegovina.
- Savva JV, Vaganov EA, 2006. Genetic and Environmental Effects Assessment in Scots Pine Provenances Planted in Central Siberia. *Mitig Adapt Strat Glob Change* 11: 269-290. <https://doi.org/10.1007/s11027-006-1026-2>.
- Stefanović V, Milanović S, Međedović S, Pintarić K, Rončević S, Sisojević D, 1980. Ekotipovi bijelog bora (*Pinus sylvestris* L.) u Bosni. Spec.ed. no 13. Faculty of Forestry and Forestry Institute, Sarajevo, Bosnia and Herzegovina, pp 9-25.
- Taeger S, Zang C, Liesebach M, Schneck V, Menzel A, 2013. Impact of climate and drought events on the growth of Scots pine (*Pinus sylvestris* L.) provenances. *Forest Ecol Manag* 307: 30-42. <https://doi.org/10.1016/j.foreco.2013.06.053>.
- Villani F, Eriksson G, 2006. Conservation and management of European chestnut (*Castanea sativa* L.) genetic resources: outputs of the CASCADE project. EUFORGEN Noble Hardwoods Network, Report of the sixth (9-11 June 2002, Alter do Chao, Portugal) and seventh meetings (22-24 April 2004, Arezzo - Italy), International Plant Genetic Resources Institute, Rome, Italy.
- Zhelev P, Lust N, 1999. Provenance study of Scots pine (*Pinus sylvestris* L.) in Belgium. I. Evaluation of phenotypical traits. *Silva Gandavensis* 64: 24-36.